# THINK TALK TREAT MIGRAINE® ALGORITHM



**AMONG A SURVEY OF** ADULTS IMPACTED BY MIGRAINE, **APPROXIMATELY...** 

**50%** 

**HESITATED TO SEEK CARE.**1

#### **REASONS FOR HESITATING INCLUDED:**



**WANTED TO MANAGE THE** SYMPTOMS ON THEIR OWN



WERE CONCERNED ABOUT NOT **BEING TAKEN SERIOUSLY** 



THOUGHT THEIR MIGRAINE **WAS NOT SERIOUS ENOUGH TO** WARRANT CONSULTING.1

THE THINK TALK TREAT MIGRAINE ALGORITHM CAN HELP YOU IDENTIFY KEY SYMPTOMS OF MIGRAINE AND DETERMINE A PROPER TREATMENT PLAN.

#### THINK TALK TREAT MIGRAINE® ALGORITHM<sup>2,3,4,5</sup>



## **THINK MIGRAINE**

Identify the 4 key signs and symptoms:

- 1. Headache
- 2. Impact
- 3. Photophobia
- 4. Nausea

A headache accompanied by ≥ 2 symptoms = 93% positive predictive value for migraine<sup>2</sup>



## **TALK MIGRAINE**

headache impact



## TREAT MIGRAINE

Use your patient's answer treatment path:

- 2 1 day/month = acute
- > 4 days/month = acute + preventive treatment



For more information on the Think Talk Treat Algorithm and additional resources to help discuss and treat migraine with your patients, visit **ThinkMigraine.com**.

- Shapiro RE, Nicholson RA, Zagar AJ, et al. Reasons for hesitating to consult for migraine care: Results of the OVERCOME (US) Study [abstract]. Headache. 2021;61:10.

  Lipton RB, Dodick D, Sadovsky R, et al. A self-administered screener for migraine in primary care. The ID MigraineTM validation study. Neurology. 2003;61:375-382.

  Ailani J, Burch RC, Robbins M. The American Headache Society consensus statement on integrating new migraine treatments into clinical practice. Headache. 2021;61:1021-1039.

  Katsarava Z, Buse D, Manack A, et al. Defining the differences between episodic migraine and chronic migraine. Curr Pain Headache Rep. 2012;16:86-92.

  Blumenfeld AM, Payne KA, Varon SF, et al. Disability, HRQQL, and resource use amongst chronic and episodic migraineurs. Results from the International Burden of Migraine Study (IBMS). Cephalalgia. 2011;31(3):301-315.